

Modifications to the breast cancer surveillance recommendations formulated in 2013 versus 2020

2013	2020
Who needs breast cancer surveillance?	
Strong recommendation to perform breast cancer surveillance for survivors treated with ≥ 20 Gy chest radiation	Strong recommendation to perform breast cancer surveillance for survivors treated with ≥ 10 Gy chest radiation
Moderate recommendation to perform breast cancer surveillance for survivors treated with 10-19 Gy chest radiation	
Weak recommendation to perform breast cancer surveillance for survivors treated with 1-9 Gy chest radiation	No recommendation included for routine breast cancer surveillance for survivors treated with 1-9 Gy chest radiation
No recommendation included for upper abdominal radiation exposing breast tissue	Moderate recommendation to perform breast cancer surveillance for survivors treated with upper abdominal radiation exposing breast tissue at a young age
No recommendation included for anthracyclines	No recommendation can be formulated for routine breast cancer surveillance for survivors treated with all types of anthracyclines in the absence of chest radiation
At what frequency should breast cancer surveillance be performed?	
Strong recommendation to perform annual breast cancer surveillance for at least up to 50 years of age	Strong recommendation to perform annual breast cancer surveillance for at least up to 60 years of age
At what age should continuation of intensive breast cancer surveillance be stopped?	
Moderate recommendation to continuation of breast cancer for survivors who are older than age 50 years	Moderate recommendations to continuation of breast cancer for survivors who are older than age 60 years
What surveillance modality should be used?	
Strong recommendation to perform mammography or breast MRI or a combination	Strong recommendation to perform mammography and breast MRI